

Advanced Trauma Modules interchange with blank modules on adult extrication manikins for realism in trauma life support and life saving first aid scenarios.

Module set includes:

Trauma intubation head	<ul style="list-style-type: none"> • Impaled object in the cheek, avulsed ear, unequal pupils, nasal bleeding, facial contusions, broken teeth and multiple lacerations • Manually generated carotid pulse • Articulates to manikin
Seat belt contusion	<ul style="list-style-type: none"> • Contusion with compound fracture clavicle from seat belt injury • Lay-on for chest
Burn arm with 1st, 2nd and 3rd degree burns	<ul style="list-style-type: none"> • Burns illustrated with blistering in progressive sequence. • Left forearm and hand articulates to manikin at the elbow.
Compound fracture radius	<ul style="list-style-type: none"> • Exposed proximal portion of radius. • May be interchanged with blank module in manikin's forearm.
Industrial hand	<ul style="list-style-type: none"> • Severe laceration to the dorsum of the hand with exposed bone and soft tissue. • Open and closed fracture of index finger and severe tear of the fingernail with contusion. • Right forearm and hand articulates to manikin at the elbow.
Exposed viscera	<ul style="list-style-type: none"> • Abdominal wound with protruding abdominal contents. • May be interchanged with blank abdominal module.
Large and small	<ul style="list-style-type: none"> • Entry wounds may be interchanged with blank

caliber entry and exit wounds	<p>deltoid pads.</p> <ul style="list-style-type: none"> • Exit wounds may be interchanged with blank ventral gluteal (small caliber) and dorsal gluteal (large caliber) pads.
Impaled Object	<ul style="list-style-type: none"> • Metallic object imbedded in thigh. • This may be interchanged with blank module in manikin's thigh.
Compound fracture femur	<ul style="list-style-type: none"> • Exposed fractured femur protruding from mid-thigh. • May be interchanged with blank module in manikin's thigh.
Closed fracture tibia and fibula	<ul style="list-style-type: none"> • Palpation of fracture is possible. • Right lower leg articulates to manikin at the knee.
Contused ankle and foot	<ul style="list-style-type: none"> • Left lower leg and foot articulates to manikin at the knee.
Crushed foot	<ul style="list-style-type: none"> • Severe lacerations, exposed bone, tendon and soft tissue. • Complete amputation of the lesser toe. • Right foot may be articulated to the manikin at the ankle.